



INVESTOR IN PEOPLE

The Patent Office
 Concept House
 Cardiff Road
 Newport

South Wales

NP1 80Q 4 JUN 2004

WIPO

PCT

PRIORITY DOCUMENT


SUBMITTED OR TRANSMITTED IN
 COMPLIANCE WITH RULE 17.1(a) OR (b)

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated

26 May 2004

BEST AVAILABLE COPY

Patents Form 1/77

Patents Act 1977
(Rule 16)



16APR03 E800562-13 D00192
P01/7700 0-00-0308732.7

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference	N.88273 GWM		
2. Patent application number (The Patent Office will fill in this part)	0308732.7		
3. Full name, address and postcode of the or of each applicant (underline all surnames)	AXCESS LIMITED 44 Esplanade, St Helier Jersey Channel Islands JE1 3UQ		
Patents ADP number (if you know it)	853854800		
If the applicant is a corporate body, give the country/state of its incorporation	British body corporate		
4. Title of the invention	ABSORPTION ENHANCERS		
5. Name of your agent (if you have one)	J.A. KEMP & CO.		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	14 South Square Gray's Inn London WC1R 5JJ		
Patents ADP number (if you know it)	26001		
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day / month / year)	
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. See note (d))	Yes		

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 15

Claim(s) 3

Abstract 1

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*) 1

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature

J.A. Kemp & Co.

Date 15 April 2003

J.A. KEMP & CO.

12. Name and daytime telephone number of person to contact in the United Kingdom

G W McCluskie
020 7405 3292

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- e) Once you have filled in the form you must remember to sign and date it.
- f) For details of the fee and ways to pay please contact the Patent Office.

ABSORPTION ENHANCERS

The present invention relates to the use of an aromatic alcohol to enhance the uptake of molecules, including biologically active macromolecules, into the body, suitably across the intestinal wall from the lumen of the gut. In particular the present invention relates to novel pharmaceutical compositions comprising an active macromolecular principle to be absorbed into the body, preferably across the intestinal wall.

Hydrophilic aromatic alcohols, in particular aromatic alcohols in which the hydroxy group is not attached directly to the aromatic nucleus, such as phenoxyethanol, phenyl ethanol and benzyl alcohol, have been employed in pharmaceutical practice for many years as solvents and plasticisers, and have a low toxicity profile when administered via various routes, including the oral route. Those compounds are all liquids at room temperature, and can be readily dissolved in aqueous media.

Hydrophilic aromatic alcohols such as phenoxyethanol and related compounds including phenyl ethanol and benzyl alcohol, have a range of actions on intestinal cells, one of which is that, when present in relatively high local concentration, aromatic alcohols transiently increase the permeability of a barrier layer of intestinal cells.

It is postulated that this is due to the opening of the tight junctions between these cells creating pores through which even large molecules (macromolecules) can pass by diffusion.

Based on the finding that an increase in the permeability of a barrier layer of intestinal cells is only seen at relatively high local concentrations of hydrophilic aromatic alcohol, the applicant's research has shown that a solution of hydrophilic aromatic alcohol co-administered orally (as an elixir) with a detectable molecule produces no enhancement of uptake. It is postulated that this is because, before it reaches the absorption site (in the intestine), the hydrophilic alcohol is rapidly diluted in the gastrointestinal tract to a concentration below which it cannot exert its effect. In addition, the molecules whose uptake one is seeking to elicit will also be diluted out before the intestine is reached. It has now been found that another class of aromatic alcohols also displays characteristics of permeation enhancers. These compounds have hydroxyl groups attached directly to the aromatic nucleus and an additional substituent in the position para to the OH group, and typically display antioxidant properties, which may or may not be related to their ability to act as

permeation enhancers. Examples of this class of compounds are propyl gallate, butylated hydroxy toluene (BHT) and butylated hydroxy anisole (BHA).

Surprisingly, although these materials have been employed routinely in pharmaceutical practice for at least twenty years primarily in lipid-based

formulations, generally as antioxidants, the observation that these materials are capable of acting as permeation enhancers has never been made. This is probably because these compounds are all solids which are sparingly soluble in water, thus making it difficult to incorporate them into water-based pharmaceutical formulations in high concentrations, and also preventing them from being available in soluble form to act as enhancers at elevated concentration when the formulation is dispersed in the lumen of the intestine, or close to any other mucosal surface where permeation enhancement is required.

The use of gallate esters or specifically propyl gallate has been described in US 6,180,666 and US 5,962,522 respectively as enhancers of bioavailability of small molecules via a mechanism in which the propyl gallate inhibits the activity of cytochrome P450 (in particular CY3PA, located in the endoplasmic reticulum), thereby reducing the metabolic degradation of small molecules on their passage through intestinal cells (known as the transcellular route). Propyl gallate and other esters of gallic acid appear to be potent inhibitors of cytochrome P450, and it is claimed that sufficient propyl gallate can be introduced into a formulation to exert a significant effect without the need for solubilisation aids. The enzyme inhibitor mechanism of action described for propyl gallate, however, cannot be expected to enhance the bioavailability of macromolecules, since macromolecules are incapable of entering unaided into intestinal cells, and so would not come into contact with the endoplasmic reticulum where the enzyme is located. Furthermore, macromolecules such as peptides and proteins are far less susceptible to the action of cytochrome P450 than are small drug molecules, so that degradation by this enzyme is not a major cause of the poor bioavailability of macromolecules from the gut, or other mucosal tissues. A much greater barrier is simply the size of the molecules themselves, which prevents them from entering into or passing through the cells lining mucosal tissues unaided, where cells which line these tissues form a continuous impassable wall.

It has now been found that, surprisingly, aromatic alcohols such as propyl gallate, BHT, BHA and analogues and derivatives thereof are capable of enhancing the passage of macromolecules across mucosal barriers by increasing the physical permeability of the mucosal cells. One possible mechanism for this to occur is by

transient opening of the tight junctions in between these cells, creating channels along which the macromolecules can pass (paracellular route). An alternative mode of action is enhancement of fluid-phase pinocytosis, resulting in internalisation of bulk fluid together with macromolecules within vacuoles, which are transported from one side of the cell to the other. While yet other mechanisms still not clearly understood are also possible, it is considered unlikely that macromolecules actually gain direct access to the internal cytoplasmic compartment of the cells. It has been found that this phenomenon is concentration-dependent, and that provision of the aromatic permeation enhancer in the high concentrations increases the effect *in vivo*. Consequently, the use of solubilisation aids is advantageous for these compounds, particularly in the case of propyl gallate, to be able to enhance the bio-availability of macromolecules from mucosal tissues.

It has now also been discovered that there are certain agents, known here as solubilisation aids, which can be used to assist in solubilising these aromatic alcohol permeation enhancers, and which, furthermore, can increase their solubility, and/or rate of dissolution when exposed to aqueous media. This is clearly important if these materials are to exert their maximal effect as permeation enhancers.

The invention provides a pharmaceutical composition comprising a mixture of:

- (a) an active macromolecular principle; and
- (b) an aromatic alcohol absorption enhancer chosen from butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof, wherein the aromatic alcohol absorption enhancer is present in an amount by weight greater than or equal to that of the active macromolecular principle.

The invention further provides a pharmaceutical composition comprising a mixture of:

- (a) an active macromolecular principle; and
- (b) an aromatic alcohol absorption enhancer chosen from propyl gallate, butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof, and
- (c) a solubilisation aid capable of increasing the solubility of the aromatic alcohol absorption enhancer in aqueous media, wherein the aromatic alcohol absorption enhancer is present in an amount by weight greater than or equal to that of the active macromolecular principle.

The invention also provides the use, in a pharmaceutical composition, of an aromatic alcohol chosen from butylated hydroxy toluene, butylated hydroxy anisole

and analogues and derivatives thereof as an enhancer for the absorption of macromolecules into the body.

In a further embodiment the invention provides the use of an aromatic alcohol chosen from butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof in the manufacture of a medicament (pharmaceutical composition) containing an active macromolecular principle, in order to enhance absorption of the active macromolecular principle into the human or animal body.

The invention also provides the use, in a pharmaceutical composition, of an aromatic alcohol chosen from propyl gallate, butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof together with a solubilisation aid capable of increasing the solubility of the aromatic alcohol absorption enhancer in aqueous media as an enhancer for the absorption of macromolecules into the body.

In a further embodiment, the invention provides the use of an aromatic alcohol chosen from propyl gallate, butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof together with a solubilisation aid capable of increasing the solubility of the aromatic alcohol absorption enhancer in aqueous media in the manufacture of a medicament (pharmaceutical composition) containing an active macromolecular principle, in order to enhance absorption of the active macromolecular principle into the human or animal body.

The aromatic alcohol absorption enhancer may be propyl gallate or an analogue or a derivative thereof, and, preferably is propyl gallate. Suitable analogues and derivatives of propyl gallate include esters of gallic acid. The esters may be linear or branched chain C_{1-12} alkyl, C_{1-12} alkyloxy, C_{1-12} alkylthio or C_{2-12} alkenyl esters. The compounds are optionally substituted with halogen, linear or branched chain C_{1-12} alkyl, C_{1-12} alkyloxy, C_{1-12} alkylthio or C_{2-12} alkenyl esters. The aromatic alcohol absorption enhancer may also be chosen from BHT, BHA and analogues and derivatives thereof. Suitable analogues and derivatives of BHT or BHA include analogues and derivatives of hydroxy toluene or hydroxy anisole where the methyl group or the methoxy group linked to the aromatic ring and/or the hydrogen ortho to the hydroxyl group are replaced by linear or branched chain C_{1-12} alkyl, C_{1-12} alkyloxy, C_{1-12} alkylthio or C_{2-12} alkenyl, either unsubstituted or substituted in any position, especially by halogen atoms. Preferably, the aromatic alcohol absorption enhancer is chosen from propyl gallate, BHT and BHA.

The aromatic alcohols disclosed above which are used in pharmaceutical practice as antioxidants are included at concentrations up to 0.1% w/v of the total formulation (see entries for individual compounds in the Handbook of

Pharmaceutical Excipients, Eds Wade & Weller, The Pharmaceutical Press, London UK, 2nd edition 1994). It is generally considered that higher concentrations of the compounds give no added antioxidant benefit, and it is thus standard pharmaceutical practice to restrict the concentration of the antioxidants in formulations to no greater than 0.1%. When used as absorption enhancers according to the present invention, however, the efficacy of these compounds is concentration dependent up to a much higher level, and their proportions in a pharmaceutical formulation are much higher than previously described in the prior art.

To the applicant's knowledge, there is no suggestion in the prior art of the use of these agents as antioxidants in pharmaceutical formulations. None of these agents play any role in enhancing absorption of macromolecules by the oral route, or that these agents may be included in formulations at levels higher than is standard pharmaceutical practice for antioxidants.

For example, EP-A-0295941 discloses a formulation for oral administration in which BHA, BHT or PG may optionally be included, so that it is clear that their presence is not essential for biological efficacy of the formulation. No concentrations of these agents are specified, and the formulation is intended as a controlled-release dosage form, in marked contrast to the present invention where immediate dissolution is desirable to ensure rapid release from the capsule.

WO-A-0222158 provides compositions comprising cyclosporin (not a macromolecule) and containing BHA, BHT and PG generally as antioxidants. Although no specific concentrations of the antioxidants are given, the use of the compounds as antioxidants suggests a level of no greater than 0.1% wt.

US-A-5756450 discloses compositions comprising low molecular water insoluble compounds including water insoluble polypeptides, especially cyclopeptides such as the cyclosporins. BHA or BHT may be included as antioxidants, again in very small quantities.

US-A-5342625 again discloses compositions comprising cyclosporins. A solubilisation aid may be present to help form a microemulsion pre-concentrate of the cyclosporin. BHA or BHT may be present at low levels as antioxidants.

BHA and BHT may also be present as antioxidants in the compositions of US-A-3996355 which comprise any drug which is stable in the presence of a vegetable oil vehicle, more specifically water-sensitive drugs having a bitter taste. Macromolecules are not envisaged.

Suitable solubilisation aids include, but are not limited to, bile acids or salts such as sodium taurocholate or taurodeoxycholate, benzyl alcohol, phenyl ethanol, phenoxyethanol, transcitol or isopropanol.

The active macromolecular principles falling within the scope of the invention include all molecules capable of having a beneficial effect when absorbed into the human or animal body, especially through the intestinal wall. The beneficial effect may be, for example, therapeutic, cosmetic or preventative such as prophylactic or contraceptive. The active macromolecular principles can be of natural (biological), synthetic or semi-synthetic origin.

Macromolecules are preferably defined as molecules having a molecular weight of over 1000 Da, preferably over 2000 Da and most preferably over 3000 Da. Examples of macromolecules, including macromolecular active macromolecular principles, include:

1. Polypeptides and proteins such as insulin; calcitonin; human serum albumin; growth hormone; growth hormone releasing factors; galanin; parathyroid hormone; blood clotting proteins such as kinogen, prothombin, fibrinogen, Factor VII, Factor VIII of Factor IX; erythropoeitins and EPO mimetics; colony stimulating factors including GCSF and GMCSF; platelet-derived growth factors; epidermal growth factors; fibroblast growth factors; transforming growth factors; GLP-1; GAG; cytokines; insulin-like growth factors; bone- and cartilage-inducing factors; neurotrophic factors; interleukins including IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12; interferons including interferon gamma, interferon - 1a, interferon alphas; TNF alpha; TNF beta; TGF-beta; cholera toxin A and B fragments; E. coli enterotoxin A and B fragments; secretin; enzymes including histone deacetylase, superoxide dismutase, catalase, adenosine deaminase, thymidine kinase, cytosine deaminase, proteases, lipases, carbohydrases, nucleotidases, polymerases, kinases and phosphatases; transport or binding proteins especially those which bind and/or transport a vitamin, metal ion, amino acid or lipid or lipoprotein such as cholesterol ester transfer protein, phospholipid transfer protein, HDL binding protein; connective tissue proteins such as a collagen, elastin or fibronectin; a muscle protein such as actin, myosin, dystrophin, or mini-dystrophin; a neuronal, liver, cardiac, or adipocyte protein; a cytotoxic protein; a cytochrome; a protein which is able to cause replication, growth or differentiation of cells; a signalling molecule such as an intra-cellular signalling protein or an extracellular signalling protein (eg hormone); trophic factors such as BDNF, CNTF, NGF, IGF, GMF, aFGF, bFGF, VEGF, NT3, T3 and HARP; apolipoproteins; antibody molecules; receptors in

soluble form such as T-cell receptors and receptors for cytokines, interferons or chemokines; proteins or peptides containing antigenic epitopes and fragments; and derivatives, conjugates and sequence variants of any of the above. These and other proteins may be derived from human, plant, animal, bacterial or fungal sources, and extracted either from natural sources, prepared as recombinants by fermentation or chemically synthesised.

2. Polynucleotides such as long-chain linear or circular single-, double- or triple-stranded DNA, single-, double- or triple-stranded RNA, oligonucleotides such as antisense DNA or RNA, and analogues thereof including PNA and phosphothioate derivates. In one embodiment it is preferred that the polynucleotides used in the invention contain a CpG motif. The coding sequence of the polynucleotide may encode a therapeutic product, in particular the coding sequence may encode an extracellular protein (e.g. a secreted protein); an intracellular protein (e.g. cytosolic, nuclear or membrane protein); a protein present in the cell membrane; a blood protein, such as a clotting protein (e.g. kinogen, prothrombin, fibrinogen factor VII, factor VIII or factor IX); an enzyme, such as a catabolic, anabolic gastrointestinal, metabolic (e.g. glycolysis or Krebs cycle), or a cell signalling enzyme, an enzyme which breaks down or modifies lipids, fatty acids, glycogen, amino acids, proteins, nucleotides, polynucleotides (e.g. DNA or RNA) or carbohydrate (e.g. protease, lipase or carbohydrase), or a protein modifying enzyme, such as an enzyme that adds or takes chemical moieties from a protein (e.g. a kinase or phosphatase); a transport or binding protein (e.g. which binds and/or transports a vitamin, metal ion, amino acid or lipid, such as cholesterol ester transfer protein, phospholipid transfer protein or an HDL binding protein); a connective tissue protein (e.g. a collagen, elastin or fibronectin); a muscle protein (e.g. actin, myosin, dystrophin or mini-dystrophin); a neuronal, liver, cardiac or adipocyte protein; a cytotoxic protein; a cytochrome; a protein which is able to cause the replication, growth or differentiation of cells; a protein which aids transcription or translation of a gene or regulates transcription or translation (e.g. a transcription factor or a protein that binds a transcription factor or polymerase); a signalling molecule, such as an intracellular or extracellular signalling molecule (e.g. a hormone); an immune system protein such as an antibody, T cell receptor, MHC molecule, cytokine (e.g. IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, TNF-, TNF-, TGF-), an interferon (e.g. IFN-, IFN-, IFN-), chemokine (e.g. MIP-1, MIP-1, RANTES), an immune receptor (e.g. a receptor for a cytokine, interferon or chemokine, such as a receptor for any of the above-mentioned cytokines, interferons or chemokines) or a cell surface marker (e.g.

macrophage, T cell, B cell, NK cell or dendritic cell surfacemarker)(eg. CD 1, 2, 3, 4, 5, 6, 7, 8, 16, 18, 19, 28, 40, or 45; or a natural ligand thereof), a trophic factor (e.g. BDNF, CNTF, NGF, IGF, GMF, aFGF, bFGF, VEGF, NT3, T5, HARP) or an apolipoprotein; a tumour suppressor (e.g. p53, Rb, Rap1A, DCC or k-rev); a suicide protein (thymidine kinase or cytosine deaminase); or a gene repressor. The proteins and peptides encoded by the polynucleotides useful in the invention may be immunogenic i.e. contain an antigen specific to the activity of the protein against which antibodies are generated by the immune system.

10 The polynucleotide may have control sequences operably linked to the coding sequence. The control sequences may typically be those of any eukaryote or of a virus which infects such eukaryotes. The polynucleotide may comprise an origin of replication.

The polynucleotides may be chemically modified. This may enhance their resistance to nucleases or may enhance their ability to enter cells. For example, 15 phosphorothioate oligonucleotides may be used. Other deoxynucleotide analogs include methylphosphonates, phosphoramidates, phosphorodithioates, N3'P5'-phosphoramidates and oligoribonucleotide phosphorothioates and their 2'-O-alkyl analogs and 2'-O-methylribonucleotide methylphosphonates. Alternatively mixed backbone oligonucleotides (MBOs) may be used. MBOs contain segments of 20 phosphothioate oligodeoxynucleotides and appropriately placed segments of modified oligodeoxy- or oligoribonucleotides. MBOs have segments of phosphorothioate linkages and other segments of other modified oligonucleotides, such as methylphosphonate, which is non-ionic, and very resistant to nucleases or 2'-O-alkyloligoribonucleotides.

25 The polynucleotide suitable for use in the invention is preferably in a form in which it is substantially free of or associated with cells or with cellular, prokaryotic, eukaryotic, nuclear, chromatin, histone or protein material. It may be in substantially isolated form, or it may be in substantially purified form, in which case it will generally comprise more than 90%, e.g. (more than or at least) 95%, 98% or 99% of 30 the polynucleotide or dry mass in the preparation. Thus the polynucleotide may be in the form of 'naked DNA'.

3. Polysaccharides such as heparin, low-molecular weight heparin, polymannose, cyclodextrins and lipopolysaccharide.

4. Any or all of the above either separately or in combination with each 35 other (for example in the form of a heteroconjugate), or with additional agents.

In preferred embodiments of the invention the active macromolecular principle to be absorbed is selected from calcitonins, insulin, low molecular weight heparin, erythropoietin, human growth hormone and parathyroid hormone.

Depending on the nature of additional excipients employed, the pharmaceutical composition of the invention may be in liquid, solid, semi-solid or gel form. The pharmaceutical composition of the invention is suitable for administration via any route giving access to different mucosal tissues such as buccal and sublingual mucosa, the nasal palate, the lungs, the rectum, the intestinal tract (including the large and small intestines) and the vagina. In the case of liquid, semi-
10 solid or gel formulations, these may be either anhydrous or aqueous.

Where the intended site of action of the composition of the invention is the intestine, it is desirable that the composition is enclosed within an enteric coating which can withstand the stomach, so that the components of the formulation remain together, undiluted and in close association until they reach the tissues of the small
15 intestine or colon. Such formulations will suitably be anhydrous. Compositions in liquid form will suitably be administered as enteric-coated capsules, while solid formulations may be administered either within enteric-coated capsules, or in tablet form.

The enteric coating is chosen appropriately to withstand the natural condition
20 of the stomach and to become permeable at the desired location in the intestine. This is preferably determined by the pH conditions which modulate along the length of the intestine. Where the site of action is the small intestine, it is preferred that the enteric coating becomes permeable and releases its contents at a pH from 3 to 7, preferably from 5.5 to 7, more preferably from 5.5 to 6.5. Where the intended site of
25 action is the colon, it is preferred that the enteric coating becomes permeable and releases its contents at a pH of 6.8 or above.

Suitable enteric coatings are well known in the art and include polymethacrylates such as those selected from the L and S series of Eudragits in particular Eudragits L12.5P, L12.5, L100, L100-55, L30 D-55, S12.5P, S12.5 and
30 S100. Selection of an appropriate coating for the capsule, which is preferably a gelatine capsule, can readily be made by the person skilled in the art based on their knowledge and the available literature supporting the Eudragit products.

Where the intended site of action is the nasal mucosa, the formulation may be in the form of an aqueous solution or as a dry powder, which can be administered as
35 a spray.

Where the intended site of action is the rectum, an appropriate method of administration is as an anhydrous liquid or solid enclosed within a capsular shell, or incorporated into the matrix of an erodible suppository.

For vaginal application, administration of the formulation in gel form is also appropriate.

In the compositions of the invention, the aromatic alcohol absorption enhancer is present in an amount (by weight) greater than or equal to that of the active macromolecular principle. This provides an effective concentration of aromatic alcohol absorption enhancer at the intestinal cell barrier layer (intestinal wall) so as to cause enhanced absorption in the co-presence of a suitable amount of the active macromolecular principle which, when absorbed, will exert its normal beneficial effect. The practitioner of the invention would select the amounts of the aromatic alcohol absorption enhancer and active macromolecular principle on the basis of the amount (for example, blood concentration level) of the active macromolecular principle concerned which is necessary for therapeutic efficacy. The weight ratio of aromatic alcohol absorption enhancer to active macromolecular principle in the mixture contained in the capsule is suitably at least 1:1, preferably at least 5:1, for example from 1:1 to 100:1, preferably from 3:1 to 50:1, most preferably from 5:1 to 20:1.

The ratio of solubilisation aid to aromatic alcohol absorption enhancer is suitably at least 1:1, preferably from 1:1 to 10:1, and most preferably from 1.5:1 to 5:1.

The absolute amount of the active macromolecular principle would be selected on the basis of the dosage of the substance required to exert the normal beneficial effect with respect to the dosage regimen used and the patient concerned. Determination of these amounts falls within the mantle of the practitioner of the invention.

In the composition for oral administration it is preferred that the contents of the capsule comprises a suitable amount of the active macromolecular principle to achieve its normal therapeutic effect. For example, the composition may contain from 0.05 to 50%, preferably from 0.1 to 25%, more preferably from 0.1 to 10% by weight of the active macromolecular principle based on the weight of the capsule contents (not including the capsule itself).

The composition of the invention may further comprise one or more other absorption enhancer compounds, for example, medium chain fatty acids and medium chain monoglycerides.

The composition of the invention may optionally further comprise any conventional additive used in the formulation of pharmaceutical products including, for example, anti-oxidants, anti-microbials, suspending agents, fillers, diluents, disintegrants, swelling agents, viscosity regulators, plasticisers and acidity regulators (particularly those adjusting the intestinal milieu to between 7 and 7.5).

The composition of the invention may optionally further comprise additional active principles which may enhance the desired action of the composition in a synergistic fashion. For example, where the active macromolecular principle is insulin, the composition may also comprise an insulin sensitiser capable of increasing the body's response to the insulin absorbed. Examples of sensitisers which could be employed in this fashion are troglitazone, pioglitazone, rosiglitazone and other members of the glitazone class of molecules.

In the composition of the invention where the mixture is contained in a capsule or tablet which comprises the aromatic alcohol absorption enhancer and active macromolecular principle, the formulation is preferably substantially anhydrous. In more preferred embodiments of the invention the entire composition is substantially anhydrous. Substantially anhydrous in the context of this invention means less than 5%, preferably less than 1% and more preferably less than 0.5% water by weight of the mixture.

The compositions of the invention can, depending on the active macromolecular principle used therein, be used in the treatment of a variety of conditions and diseases of the human or animal body by therapy or, alternately, can be used to introduce macromolecules essential for the diagnosis of diseases and conditions within the human or animal body. The compositions of the invention are preferably pharmaceutical or cosmetic compositions.

In the compositions of the invention the mixture contained in the capsule may be a liquid, semi-solid or gel, which is either in the form of a solution or a microparticulate dispersion. That is to say the active macromolecular principle(s) for absorption are incorporated into the formulation either in the form of a solution or as a microparticulate dispersion. Alternatively, the composition may be in the form of a solid.

The compositions of the invention are suitably produced by preparing a substantially anhydrous mixture of the active macromolecular principle and the aromatic alcohol absorption enhancer and then optionally filling uncoated capsules with the mixture and optionally coating them with an appropriate polymer mixture to achieve the desired permeability properties.

The following Examples serve to illustrate the present invention and should not be construed as limiting.

EXAMPLES

Example 1 Effect in permeabilising cell culture monolayer

Caco-2 cells (a cell line derived from human colon adenocarcinoma) are grown as a confluent mono-layer on the surface of a porous membrane (pore size $0.4\mu\text{m}$, surface area 0.33cm^2) separating two aqueous compartments, the upper compartment filled with $200\mu\text{l}$ of culture medium, and the lower compartment containing $600\mu\text{l}$ of the culture medium. Electrical resistance across the mono-layer is measured using an epithelial volttohmmeter connected to electrodes inserted into the medium on either side of the mono-layer in the upper and lower compartments. This trans-epithelial electrical resistance (TEER) is measured immediately before, and fifteen minutes after the addition of aromatic alcohols to the upper compartment (typical results are given in the table below). Four replicates are employed for each compound, whose concentrations are shown in the table below. Fall in TEER is considered indicative of the increased flow of materials (including bulk fluid phase) across the cell monolayer.

Falls of greater than 50% of the initial value are considered significant. Reducing the concentration tends to reduce the effect observed.

Table

Agent	Concentration (mg/ml)	TEER (ohm.cm^2)	
		Before addition	15 minutes after addition
Propyl gallate	13	494	-0.6
Propyl gallate	0.2	667	629
Butylated hydroxy anisole	0.3	564	183
Butylated hydroxy toluene	0.3	633	194

Example 2 Preparation of formulation containing insulin, propyl gallate and sodium taurocholate

Sodium taurocholate in an amount of 150mg is mixed with 75mg of propyl gallate in a glass vial and $825\mu\text{l}$ of distilled water are added. Dissolution at room temperature is not achieved even on prolonged shaking, but after warming with brief sonication in an ultrasonic bath a clear colourless solution is obtained. Bovine

insulin in an amount of 8.4mg is added to the solution with mixing, followed by 10 μ l of glacial acetic acid while vortexing the insulin suspension. A clear solution is rapidly obtained, with a pH of 3.15. The contents of the vial are frozen rapidly with shaking and lyophilised overnight. The following day a dry solid is obtained. An amount of 10mg of the solid is weighed into a 2ml vial and 50 μ l of distilled water added. A clear solution forms rapidly.

Example 3 Preparation of formulation containing insulin, propyl gallate and sodium taurodeoxycholate

Identical conditions to those described in example 2 are employed using taurodeoxycholate instead of taurocholate. The pH of the final solution before drying is 3.36. A clear solution forms rapidly on addition of distilled water to the dried solid as before.

Example 4 Preparation of formulation containing calcitonin, propyl gallate and sodium taurocholate

Identical conditions to those described in example 2 are employed, except that 2.3mg of salmon calcitonin is dissolved in distilled water, and the entire solution added to the mixture of sodium taurocholate and propyl gallate. A clear solution forms rapidly on addition of distilled water as before.

Example 5 Preparation of formulation containing calcitonin, propyl gallate and sodium taurodeoxycholate

Identical conditions to those described in example 4 are employed, except that taurodeoxycholate is employed instead of taurocholate.

Example 6 Preparation of formulation containing parathyroid hormone, propyl gallate and sodium taurocholate

Identical conditions to those described in example 4 are employed, except that 0.5mg parathyroid hormone is employed instead of calcitonin.

Example 7 Preparation of formulation containing parathyroid hormone, propyl gallate and sodium sodium taurodeoxycholate

Identical conditions to those described in example 6 are employed, except that taurodeoxycholate is employed instead of taurocholate.

Example 8 Preparation of formulation containing parathyroid hormone, propyl gallate and sodium taurodeoxycholate

Identical conditions to those described in example 7 are employed, except that the bile salt/PG mixture is dried without addition of protein, and parathyroid hormone is added as a dry powder to the dry residue after lyophilisation.

Example 9 Preparation of formulation containing human growth hormone, propyl gallate and sodium taurodeoxycholate

Identical conditions to those described in example 8 are employed, except that 20mg of human growth hormone is employed instead of parathyroid hormone.

Example 10 Preparation of formulation containing calcitonin, propyl gallate and propylene glycol

75mg of propyl gallate is dissolved by vortexing in 200µl propylene glycol. 200µl of the resultant solution is then transferred to a vial containing 1mg of solid calcitonin. The vial is vortexed briefly to disperse the solid, then shaken for one hour at 37°C, giving a clear solution.

Example 10 Preparation of formulation containing calcitonin, propyl gallate and benzyl alcohol

100mg of propyl gallate is vortexed in 200µl of benzyl alcohol, giving a clear solution after several minutes at room temperature. 200µl of the resultant solution is then transferred to a vial containing 1mg of solid calcitonin. The vial is vortexed briefly to disperse the solid.

Example 11 Preparation of formulation containing calcitonin, propyl gallate and transcutool

100mg of propyl gallate is vortexed in 200µl of transcutool, giving a clear solution after one minute at room temperature. 200µl of the resultant solution is then transferred to a vial containing 1mg of solid calcitonin. The vial is vortexed briefly to disperse the solid for one hour at 37°C, giving a clear solution. 200µl of the resultant solution is then transferred to a vial containing 1mg of solid calcitonin. The vial is vortexed briefly to disperse the solid, then shaken for one hour at 37°C, giving a clear solution. 100µl of the solution is transferred to a fresh vial to which 100µl of distilled water is added. All components remain in solution as a single-phase clear liquid.

Example 12 Preparation of formulation containing calcitonin, butylated hydroxytoluene and transcitol

100mg of butylated hydroxy toluene is vortexed in 200 μ l of transcitol, giving a clear solution after several minutes at room temperature. 200 μ l of the
5 resultant solution is then transferred to a vial containing 1mg of solid calcitonin. The vial is vortexed briefly to disperse the solid, then shaken for one hour at 37°C, giving a clear solution. 100 μ l of the solution is transferred to a fresh vial to which 100 μ l of distilled water is added, giving a clear opalescent solution at 37°C.

CLAIMS

1. A pharmaceutical composition comprising a mixture of :
 - (a) an active macromolecular principle, and
 - (b) an aromatic alcohol absorption enhancer chosen from butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof, wherein the aromatic alcohol absorption enhancer is present in an amount by weight greater than or equal to that of the active macromolecular principle.
2. A pharmaceutical composition comprising a mixture of :
 - (a) an active macromolecular principle,
 - (b) an aromatic alcohol absorption enhancer chosen from propyl gallate, butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof, wherein the aromatic alcohol absorption enhancer is present in an amount by weight greater than or equal to that of the active macromolecular principle, and
 - (c) a solubilisation aid capable of increasing the solubility of the aromatic alcohol absorption enhancer in aqueous media.
3. A composition according to claim 1 or 2, wherein the mixture comprises less than 5% by weight of water.
4. A composition according to any of claims 1 to 3, wherein the composition is coated with an enteric coating which becomes permeable at a pH of from 3 to 7.
5. A composition according to any preceding claim, wherein the mixture comprises at least 1% by weight of the aromatic alcohol absorption enhancer.
6. A composition according to any preceding claim, wherein the ratio by weight of the aromatic alcohol absorption enhancer to active macromolecular principle is at least 5:1.
7. A composition according to any preceding claim, wherein the mixture is in the form of a solution or a microparticulate dispersion.
8. A composition according to any preceding claim, wherein the mixture is in solid form.
9. A composition according to any preceding claim, wherein the active macromolecular principle is a polypeptide or protein, polynucleotide, polysaccharide or a mixture thereof.
10. A composition according to any preceding claim, wherein the aromatic alcohol absorption enhancer is chosen from BHT, BHA and analogues and

derivatives thereof, including analogues and derivatives of hydroxy toluene or hydroxy anisole where the methyl group or the methoxy group linked to the aromatic ring and/or the hydrogen ortho to the hydroxyl group are replaced by linear or branched chain C₁₋₁₂ alkyl, C₁₋₁₂ alkyloxy, C₁₋₁₂ alkylthio or C₂₋₁₂ alkenyl, either
 5 unsubstituted or substituted in any position, especially by halogen atoms.

11. A composition according to any one of claims 2 to 9, wherein the aromatic alcohol absorption enhancer is propyl gallate or an analogue or a derivative thereof, including esters of gallic acid, where the esters may be linear or branched chain C₁₋₁₂ alkyl, C₁₋₁₂ alkyloxy, C₁₋₁₂ alkylthio or C₂₋₁₂ alkenyl esters, and the
 10 compounds are optionally substituted with halogen, linear or branched chain C₁₋₁₂ alkyl, C₁₋₁₂ alkyloxy, C₁₋₁₂ alkylthio or C₂₋₁₂ alkenyl esters.

12 A composition according to any of claims 2 to 11, where the solubilisation aid is chosen from a bile acid or salt, benzyl alcohol, phenyl ethanol, phenoxyethanol, transcitol and isopropanol.

13. A composition according to any preceding claim, where the active macromolecular principle is insulin, calcitonin, growth hormone, parathyroid hormone, or erythropoietin, and derivatives and analogues, either synthetic or from
 15 natural sources, conforming to structures derived from either human or animal origin.

14. A composition according to any preceding claim, where the active macromolecular principle is insulin or a derivative or an analogue thereof, either
 20 synthetic or from natural sources, conforming to structures derived from either human or animal origin, and the composition further comprises an insulin sensitising agent.

15. A composition according to any preceding claim, for use in the therapeutic or diagnostic treatment of the human or animal body.
 25

16. Use, in a pharmaceutical composition, of an aromatic alcohol chosen from butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof as an enhancer for the absorption of a macromolecule across the intestinal wall.

17. Use of an aromatic alcohol chosen from butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof in the manufacture of a medicament containing an active macromolecular principle, in order to enhance
 30 absorption of the active macromolecular principle into the human or animal body.

18. Use, in a pharmaceutical composition, of an aromatic alcohol chosen from propyl gallate, butylated hydroxy toluene, butylated hydroxy anisole and
 35 analogues and derivatives thereof together with a solubilisation aid capable of

increasing the solubility of the aromatic alcohol absorption enhancer in aqueous media as an enhancer for the absorption of macromolecules across the intestinal wall.

19. Use of an aromatic alcohol chosen from propyl gallate, butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof

5 together with a solubilisation aid capable of increasing the solubility of the aromatic alcohol absorption enhancer in aqueous media in the manufacture of a medicament containing an active macromolecular principle, in order to enhance absorption of the active macromolecular principle into the human or animal body.

20. Use according to any one of claims 16 to 19, wherein the composition
10 comprises less than 5% by weight of water.

21. Use according to any of claims 18 or 19, wherein the solubilisation aid is chosen from a conjugated bile acid or salt, benzylalcohol, phenylethanol, phenoxyethanol, transcutool and isopropanol.

22. Use according to any of claims 17 or 19, wherein the medicament is
15 provided in the form of a solution, as a microparticulate dispersion or as a solid.

23. Use according to any of claims 16 to 22, wherein the macromolecules to be absorbed/active macromolecular principle is a polypeptide or protein, polynucleotide, polysaccharide or a mixture thereof.

24. Use according to claim 23, wherein the macromolecules to be
20 absorbed/active macromolecular principle to be absorbed is chosen from insulin, calcitonin, growth hormone, parathyroid hormone and erythropoietin, and derivatives and analogues thereof, either synthetic or from natural sources, conforming to structures derived from either human or animal origin.

25. Use according to claim 24, wherein the macromolecules to be
25 absorbed/active macromolecular principle to be absorbed is insulin or a derivative or an analogue thereof, either synthetic or from natural sources, conforming to structures derived from either human or animal origin, and the medicament further comprises an insulin sensitising agent.

26. A method of enhancing the absorption of an active macromolecular
30 principle in a patient, which method comprises administering to said patient a composition as defined in any one of claims 1 to 14.

27. A method of treating a patient suffering from a condition or disease treatable by administration of a composition according to any of claims 1 to 14.

ABSTRACT

The invention provides a pharmaceutical composition comprising a mixture of:

- 5 (a) an active macromolecular principle, and
 (b) an aromatic alcohol absorption enhancer chosen from butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof, wherein the aromatic alcohol absorption enhancer is present in an amount by weight greater than or equal to that of the active macromolecular principle, and further
- 10 comprises a pharmaceutical composition comprising a mixture of :
- (a) an active macromolecular principle,
 (b) an aromatic alcohol absorption enhancer chosen from propyl gallate, butylated hydroxy toluene, butylated hydroxy anisole and analogues and derivatives thereof, wherein the aromatic alcohol absorption enhancer is present in an amount by
- 15 weight greater than or equal to that of the active macromolecular principle, and
 (c) a solubilisation aid capable of increasing the solubility of the aromatic alcohol absorption enhancer in aqueous media.

PCT/GB2004/001650



This Page is inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLORED OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images
problems checked, please do not report the
problems to the IFW Image Problem Mailbox**